

Copyright is owned by the Author of the thesis. Permission is given for a copy to be downloaded by an individual for the purpose of research and private study only. The thesis may not be reproduced elsewhere without the permission of the Author.

Anatomical studies of claw conformation in New Zealand dairy cattle

A thesis submitted in partial fulfilment of the requirements for the degree of

Doctor of Philosophy

in

Veterinary Science

Massey University, Palmerston North, New Zealand

Linda Jean Laven

2014

Abstract

The aim of the studies reported in this thesis was to extend our current knowledge on estimating and monitoring claw conformation under New Zealand conditions.

Initially post mortem material was used to: (1) evaluate the utility of claw volume estimation with data from cull dairy cows; (2) validate the use of a portable ultrasound machine in the estimation of internal claw dimensions; and (3) apply morphometry to quantify vascular change in histological sections from the collected claw material.

Analysis of the relationship between claw volume and hoof conformation revealed differences between the claws of dairy cattle used for this study and those of beef cattle which had been used previously. This necessitated the development of a different predictive model for dairy cattle which was subsequently used in the live animal studies. Validation of ultrasound estimates for sole and soft tissue thickness against calliper measurements in the sectioned claw, found that the portable machine used accurately estimated mean distance to distal phalanx (DP) and was thus suitable for categorising claws as having thin, marginal or adequate sole thickness.

Morphometrical techniques were successfully applied to sections from frozen claw material; the ratio of vessel lumen area to overall vessel area was found to vary depending on site, claw and overall vessel size. No relationship was found between the ratio and claw horn haemorrhages, but the level of such lesions was low in the animals available for analysis.

Subsequently, selected conformation traits were assessed on-farm in two consecutive cohorts of first lactation heifers. Some variables were identified as being useful to evaluate conformational change within a pasture-based system. These demonstrated dynamic change over the course of lactation. However, fluctuation in these variables at pasture means that any intervention study would need to have a greater impact on the claw than the variations produced by the background environment. Other variables showed stability over lactation and between years of study, and are therefore potentially suited to the investigation of claw size in relation to the development of lesions and lameness.

In Year 1, ultrasound estimates of DP were recorded as a proxy for sole thickness. Values decreased significantly after calving to a nadir at approximately Day 110. The change in DP between Days 10 and 110 was associated with the initial value of DP on Day 10. The study concluded that thin soles increased in depth while thick soles wore to become thinner, indicating that heifers with thinner soles were able to accommodate to the changes occurring around and after calving and that the response of the claw to encountering tracks, collecting yards and milking parlours is not simply an increase in net wear.

In Year 2, a novel method to capture changes in heel conformation was successfully trialled. Non-weight bearing heel length was found to alter rapidly after calving while other measures responded slowly in comparison and suggested extended monitoring was advisable. The on-farm studies established that, for most conformational variables, there is value in recording information from all four claws of the hind feet in a pasture based system.

Acknowledgements

Thank you to everyone who helped and encouraged me to complete this thesis.

There are so many people who said the right words at the right time to keep me going and for that please know I am very grateful. I especially mention my immediate family who have lived and grown around this work, my extended family, friends and colleagues for their support and advice. To delegates at the International Ruminant Lameness Symposium, Kuopio (Finland), Rotorua (New Zealand) and Bristol (UK), thank you for your feedback and enthusiasm for all aspects of lameness.

Special thanks go to my supervision panel: Professor T.J. Parkinson, Professor N. Lopez-Villalobos, Associate Professor R.A. Laven and Dr. J. K. Margerison.

In addition, I have received wonderful technical support over the years, which has been integral to the process of producing this thesis. So a massive thank you to K. Fraser, R.N. Ward, A. Nutman and M. Hogan, for help with equipment (the hoof photography box!), band saw training, and collection of material for the post mortem based studies.

Dr. M. Perrott and the histology team, for their assistance and perseverance with regard to the production of the histological sections and to K. Fraser and A. Robarts for their on-farm technical assistance, particularly the early mornings devoted to drafting.

I would like to acknowledge financial assistance for the post mortem based studies from the McGeorge research fund and support for the on-farm studies from Bell-Booth Ltd. Palmerston North New Zealand.

Contents

Anatomical studies of claw conformation in New Zealand dairy cattle.....	1
Abstract.....	i
Acknowledgements.....	ii
List of Figures	ix
List of Tables	xvi
Chapter one: Literature review.....	1
Introduction	1
Toe Angle	4
Method of assessment.....	4
Cow effects on toe angle	9
Management effects on toe angle.....	13
Links to other conformational parameters, lameness, lesions and locomotion.....	19
Dorsal border length, toe length	20
Method of assessment.....	20
Cow effects on dorsal border length	24
Management effects on dorsal border length	26
Links to other conformational parameters, lameness, lesions and locomotion.....	29
Heel conformation	30
Heel height (heel depth)	30
Method of assessment.....	30
Cow effects on heel height	33
Management effects on heel height	35
Links to other conformational parameters, lameness, lesions and locomotion.....	38
Heel Angle	40

Bulb or heel length.....	42
Claw Diagonal.....	44
Claw length (sole length, claw diagonal length, sole diagonal length).....	46
Toe height (claw height)	50
Heel width (sole breadth, claw width).....	51
Hoof base area	55
Abaxial groove height	55
Ground surface area of the claw	55
Introduction to experimental chapters	57
Overarching Hypothesis.....	59
References	59
Chapter two: Predicting the volume of the bovine claw – method validation and development.....	64
Introduction	64
Materials and methods.....	67
Experiment 1: Volumetric modelling.....	67
Experiment 2: Investigation of the relationship between predicted volume and calliper measurements of sole thickness and soft tissue depth.	70
Statistical analysis.	71
Experiment 2: Investigation of the relationship between predicted volume and calliper measurements of sole thickness and soft tissue depth.	73
Results.....	74
Water displacement.....	74
Experiment 1: Volumetric modelling - Whole foot volume modelling procedures	74
Experiment 1: Volumetric modelling - Individual claw modelling.....	86

Experiment 2: Investigation the relationship between predicted volume and calliper measurements of sole thickness and soft tissue depth.	95
Discussion.....	98
Conclusion.....	102
References	102
Chapter three: Validation of a portable ultrasound machine for the assessment of sole thickness in the New Zealand dairy cow.....	104
Introduction	104
Materials and methods.....	107
Ultrasound assessment of claws.....	108
Calliper measurement of distance to distal phalanx and sole thickness.....	108
Assessment of dorsal border and toe angle relationship to sole thickness.....	109
Statistical analysis	109
Results.....	110
Ultrasound assessment of claws.....	110
Relationship between sole thickness and dorsal border, toe angle at Sites 1 and 2.....	118
Discussion.....	120
Conclusion.....	124
Publication	125
References	125
Chapter four: Histological preparation pilot study.....	127
Introduction	127
Materials and methods.....	129
Effect of altering cryosectioning method (Protocols 1-4)	132
Paraffin block mounting vs. fast cryosectioning (Protocols 5 and 6)	132

Results.....	133
Cutting on the band saw.....	133
Cryosection mounting.....	133
Effect of freezing speed	133
Fresh versus frozen claw tissue.	134
Discussion.....	134
Conclusion.....	136
References	136
Chapter five: Morphometric evaluation of the vasculature in light microscopy sections of the bovine claw	138
Introduction	138
Materials and methods.....	141
Sample preparation	141
External sole and white line lesions.....	143
Morphometry.....	146
Statistical analysis	148
Results.....	153
Model 1: The effect of foot and site on the ratio of lumen area to total vessel area (L/V).	153
Model 2: The effect of hoof conformation on SQRT L/V.....	153
The effect of vessel size on SQRT L/V.	154
Model 3: The effect of foot and site on SQRT L/V with log ₁₀ lumen as a covariate.	155
Model 4: The effect of hoof conformation on SQRT L/V with log ₁₀ lumen as a covariate.	155
Model 5: The effect of lesion score category for the white line and sole haemorrhages on SQRT L/V.	156

Discussion.....	156
Conclusion.....	161
Further research	161
Publication	161
References	162
Chapter six: Estimations of claw conformation and volume change over lactation for two sequential cohorts of first lactation heifers.....	165
Introduction	165
Materials and Methods.....	167
Animals.....	167
Conformational measurements.....	167
Volume estimation.....	168
Statistical analysis	168
Results.....	169
Toe angle Year1.....	170
Toe angle Year 2.....	172
Dorsal border Year 1	174
Dorsal border Year 2	176
Abaxial groove Year1	178
Abaxial groove Year 2	180
Proximal: Length of coronary band from flexure of the dorsal border to the abaxial groove Year 1	182
Proximal: Length of coronary band from flexure of the dorsal border to the abaxial groove Year 2	184
Distal: Length of abaxial wall from the apex of the toe to the abaxial groove at ground surface Year 1.....	186

Distal: Length of abaxial wall from the apex of the toe to the abaxial groove at ground surface Year 2.....	188
Change in predicted claw volume over lactation for two sequential cohorts of first lactation heifers Year 1	190
Change in predicted claw volume over lactation for two sequential cohorts of first lactation heifers Year 2	192
Change in variables over time: last estimate minus first estimate for each variable ...	194
Discussion.....	196
Conclusion.....	203
References	205
Chapter seven: Live animal estimations of distance to distal phalanx and sole thickness in a cohort of twenty five first lactation heifers.....	207
Introduction	207
Materials and Methods.....	209
Animals.....	209
Ultrasound measurements	210
Ultrasound assessment of claws.....	210
Statistical analysis	211
Results.....	212
Ultrasound measurement of distance to distal phalanx	212
Ultrasound measurement of sole thickness	218
Relationship between initial depth on Day 10 and the nadir for each site.....	219
Sole thickness categories for studied claws Days 10-220.....	224
Discussion.....	229
Conclusion.....	233
Publication	234

References	234
Chapter eight: Study of heel conformation changes in first lactation heifers	236
Introduction	236
Materials and Methods.....	237
Animals.....	237
Conformational measurements.....	237
Additional heel conformation measurements.....	238
Estimation of heel and sole weight bearing surface area.	240
Statistical analysis	241
Results.....	242
Claw depth at the distal end of abaxial groove	242
Non-weight bearing heel length	244
Weight bearing heel length	246
Surface area of the weight-bearing heel	248
Sole surface area.....	250
Inter-variable relationships.....	252
Discussion.....	258
Conclusion.....	263
References	264
Chapter nine: General Discussion	266
Morphometry.....	266
Foot volume	268
Conformation variables	270
Sole thickness.....	273
Heel conformation, sole thickness and wear	274

Conclusions	279
References:	282
Appendix:	286